


Pankreatitida chronická (laboratorní diagnostika)

Chronická pankreatitida (CHP) je definována jako pokračující zánětlivé onemocnění pankreatu s charakteristickými, ireverzibilními, morfologickými změnami, způsobující charakteristické bolesti a trvalé ztráty funkce pankreatu. Laboratorní diagnostika je proto zaměřena na přímé a nepřímé funkční testy exokrinního pankreatu, stanovení míry sekrece hodnocením aktivity pankreatických enzymů – chymotrypsinu nebo elastázy-1 ve stolici. V poslední době se objevují další nové funkční testy – dechové testy, kdy perorálně podaný substrát je značen izotopem uhlíku ^{13}C (starší testy izotopem ^{14}C). Mírou aktivity pankreatických enzymů je koncentrace ^{13}C (nebo ^{14}C) ve vydechovaném vzduchu. Zlatým standardem funkční diagnostiky je sekretin-pankreozyminový (sekretin-CCK) test, jako referenční test je používáno i kvantitativní stanovení tuků v 72-hodinovém sběru stolice.



Chronická pankreatitida – zobrazovací metody

Odkazy

Související články

- Akutní pankreatitida
- Akutní pankreatitida (laboratorní diagnostika)
- Chronická pankreatitida

Zdroj

- se svolením autora převzato z KOCNA, Petr. *GastroLab : MiniEncyklopedie laboratorních metod v gastroenterologii* [online]. ©2002. Poslední revize 2011-01-08, [cit. 2011-03-04]. <<http://www1.lf1.cuni.cz/~kocna/glab/glency1.htm>>.

Použitá literatura

- BILGIN, M, et al. MRI and MRCP findings of the pancreas in patients with diabetes mellitus: compared analysis with pancreatic exocrine function determined by fecal elastase 1. *J Clin Gastroenterol.* 2009, vol. 43, no. 2, s. 165-70, ISSN 0192-0790 (Print), 1539-2031 (Electronic). PMID: 18797409 (<https://www.ncbi.nlm.nih.gov/pubmed/18797409>).
- LIEB, JG, et al. Pancreatic function testing: here to stay for the 21st century. *World J Gastroenterol.* 2008, vol. 14, no. 20, s. 3149-58, ISSN 1007-9327. PMID: 18506918 (<https://www.ncbi.nlm.nih.gov/pubmed/18506918>).
- TESTONI, PA, et al. MRCP-secretin test-guided management of idiopathic recurrent pancreatitis: long-term outcomes. *Gastrointest Endosc.* 2008, vol. 67, no. 7, s. 1028-34, ISSN 0016-5107 (Print), 1097-6779 (Electronic). PMID: 18179795 (<https://www.ncbi.nlm.nih.gov/pubmed/18179795>).
- STEVENS, T, et al. A prospective crossover study comparing secretin-stimulated endoscopic and Dreiling tube pancreatic function testing in patients evaluated for chronic pancreatitis. *Gastrointest Endosc.* 2008, vol. 67, no. 3, s. 458-66, ISSN 0016-5107 (Print), 1097-6779 (Electronic). PMID: 18294508 (<https://www.ncbi.nlm.nih.gov/pubmed/18294508>).
- GILLAMS, A, et al. Correlation of MRCP quantification (MRCPQ) with conventional non-invasive pancreatic exocrine function tests. *Abdom Imaging.* 2008, vol. 33, no. 4, s. 469-73, ISSN 0942-8925 (Print), 1432-0509 (Electronic). PMID: 17653788 (<https://www.ncbi.nlm.nih.gov/pubmed/17653788>).
- GILLAMS, AR, et al. Quantitative secretin MRCP (MRCPQ): results in 215 patients with known or suspected pancreatic pathology. *Eur Radiol.* 2007, vol. 17, no. 11, s. 2984-90, ISSN 0938-7994 (Print), 1432-1084 (Electronic). PMID: 17619882 (<https://www.ncbi.nlm.nih.gov/pubmed/17619882>).
- SCHNEIDER, AR, et al. Does secretin-stimulated MRCP predict exocrine pancreatic insufficiency?: A comparison with noninvasive exocrine pancreatic function tests. *J Clin Gastroenterol.* 2006, vol. 40, no. 9, s. 851-5, ISSN 0192-0790 (Print), 1539-2031 (Electronic). PMID: 17016144 (<https://www.ncbi.nlm.nih.gov/pubmed/17016144>).
- BALI, MA, et al. Evaluation of somatostatin inhibitory effect on pancreatic exocrine function using secretin-enhanced dynamic magnetic resonance cholangiopancreatography: a crossover, randomized, double blind, placebo-controlled study. *Pancreas.* 2006, vol. 32, no. 4, s. 346-50, ISSN 0885-3177 (Print), 1536-4828 (Electronic). PMID: 16670616 (<https://www.ncbi.nlm.nih.gov/pubmed/16670616>).
- STEVENS, T, et al. A randomized crossover study of secretin-stimulated endoscopic and dreiling tube

pancreatic function test methods in healthy subjects. *Am J Gastroenterol*. 2006, vol. 101, no. 2, s. 351-5, ISSN 0002-9270 (Print), 1572-0241 (Electronic). PMID: 16454842 (<https://www.ncbi.nlm.nih.gov/pubmed/16454842>).

- WILSON, RB, et al. Serum elastase in the diagnosis of acute pancreatitis: a prospective study. *ANZ J Surg*. 2005, vol. 75, no. 3, s. 152-6, ISSN 1445-1433 (Print), 1445-2197 (Electronic). PMID: 15777396 (<https://www.ncbi.nlm.nih.gov/pubmed/15777396>).
- AL-BAHRANI, AZ, et al. Clinical laboratory assessment of acute pancreatitis. *Clin Chim Acta*. 2005, vol. 362, no. 1-2, s. 26-48, ISSN 0009-8981 (Print), 1873-3492 (Electronic). PMID: 16024009 (<https://www.ncbi.nlm.nih.gov/pubmed/16024009>).
- PAPACHRISTOU, GI, et al. Inflammatory markers of disease severity in acute pancreatitis. *Clin Lab Med*. 2005, vol. 25, no. 1, s. 17-37, ISSN 0272-2712 (Print), 1557-9832 (Electronic). PMID: 15749230 (<https://www.ncbi.nlm.nih.gov/pubmed/15749230>).
- SIEGMUND, E, et al. The diagnostic validity of non-invasive pancreatic function tests--a meta-analysis. *Z Gastroenterol*. 2004, vol. 42, no. 10, s. 1117-28, ISSN 0044-2771 (Print), 1439-7803 (Electronic). PMID: 15508057 (<https://www.ncbi.nlm.nih.gov/pubmed/15508057>).
- KEIM, V, et al. Evaluation of Pankrin, a new serum test for diagnosis of acute pancreatitis. *Clin Chim Acta*. 2003, vol. 332, no. 1-2, s. 45-50, ISSN 0009-8981 (Print), 1873-3492 (Electronic). PMID: 12763279 (<https://www.ncbi.nlm.nih.gov/pubmed/12763279>).
- TAKEDA, M, et al. Fecal elastase-1 test: clinical evaluation of a new noninvasive pancreatic function test. *Rinsho Byori*. 2002, vol. 50, no. 9, s. 893-8, ISSN 0047-1860. PMID: 12386968 (<https://www.ncbi.nlm.nih.gov/pubmed/12386968>).
- WALKOWIAK, J, et al. Fecal elastase-1 is superior to fecal chymotrypsin in the assessment of pancreatic involvement in cystic fibrosis. *Pediatrics*. 2002, vol. 110, no. 1, s. e7-e7, ISSN 0031-4005 (Print), 1098-4275 (Electronic). PMID: 12093988 (<https://www.ncbi.nlm.nih.gov/pubmed/12093988>).
- KYLÄNPÄÄ-BÄCK, ML, et al. Comparison of urine trypsinogen-2 test strip with serum lipase in the diagnosis of acute pancreatitis. *Hepatogastroenterology*. 2002, vol. 49, no. 46, s. 1130-4, ISSN 0172-6390. PMID: 12143219 (<https://www.ncbi.nlm.nih.gov/pubmed/12143219>).
- LEMPINEM, M, et al. Predicting the severity of acute pancreatitis by rapid measurement of trypsinogen-2 in urine. *Clin Chem*. 2001, vol. 47, no. 12, s. 2103-7, ISSN 0009-9147 (Print), 1530-8561 (Electronic). PMID: 11719473 (<https://www.ncbi.nlm.nih.gov/pubmed/11719473>).
- TREACY, J, et al. Evaluation of amylase and lipase in the diagnosis of acute pancreatitis. *ANZ J Surg*. 2001, vol. 71, no. 10, s. 577-82, ISSN 1445-1433 (Print), 1445-2197 (Electronic). PMID: 11552931 (<https://www.ncbi.nlm.nih.gov/pubmed/11552931>).
- PEZZILLI, R, et al. Behaviour of serum pancreatic enzymes in chronic pancreatitis. *Dig Liver Dis*. 2000, vol. 32, no. 3, s. 233-7, ISSN 1590-8658 (Print), 1878-3562 (Electronic). PMID: 1097577 (<https://www.ncbi.nlm.nih.gov/pubmed/1097577>).