

Mikroskop:20063/annot

[

```
{ "idx": "20063+", "txt_cz": "Adventicie", "txt_en": "", "desc_cz": "", "desc_en": "", "insert_who": "Gurka", "marks": [ { "shape": "arrow", "ori": "down", "color": "green", "size": "small", "x_rel": "0.163186", "y_rel": "0.508995" } ] }, { "idx": "20063+", "txt_cz": "Lamina propria mucosae", "txt_en": "", "desc_cz": "The lamina propria\n\r\ncontains elastic fibers and is surrounded by smooth muscle", "desc_en": "", "insert_who": "Gurka", "marks": [ { "shape": "arrow", "ori": "down", "color": "green", "size": "small", "x_rel": "0.270026", "y_rel": "0.501852" } ] }, { "idx": "20063+", "txt_cz": "luman of seminal vesicle", "txt_en": "", "desc_cz": "might contain eosinophilic substnace", "desc_en": "", "insert_who": "Gurka", "marks": [ { "shape": "arrow", "ori": "down", "color": "green", "size": "small", "x_rel": "0.69566", "y_rel": "0.519962" } ] }, { "idx": "20063+", "txt_cz": "mucosa", "txt_en": "", "desc_cz": "highly convulated folds \n\r\n The folds are lined with simple (with developed golgy ap.) or pseudostratified columnar epithelial cells rich in secretory granules.\n\r\n (the pseudostratified cells have nuclei located at base)\n\r\nbasal cells - smmal , round stem cells next to the basement membrane", "desc_en": "", "insert_who": "Gurka", "marks": [ { "shape": "arrow", "ori": "down", "color": "green", "size": "small", "x_rel": "0.697922", "y_rel": "0.506138" } ] }, { "idx": "20063+", "txt_cz": "Tunica muscularis", "txt_en": "", "desc_cz": "with inner circular and outer longitudinal layers that empty the gland during ejaculation", "desc_en": "", "insert_who": "Gurka", "marks": [ { "shape": "arrow", "ori": "down", "color": "green", "size": "small", "x_rel": "0.227469", "y_rel": "0.501962" } ] }
```

```
        ],
        {
            "idx": "20063+",
            "txt_cz": "\u017dl\u00e1zov\u00fd epitel jednovrstevn\u00fd kubick\u00fd a\u017e cylindrick\u00fd",
            "txt_en": "pseudostratified epithelium",
            "desc_cz": "complex folds that fill much of the lumen .\r\n\r\nThe folds are lined with simple or pseudostratified columnar epithelial cells rich in secretory granules.",
            "desc_en": "testosteron level in blood can actually affect the size and activity of the epithelial cells in seminal vesicel.",
            "insert_who": "Gurka",
            "marks": [
                {
                    "shape": "arrow",
                    "ori": "down",
                    "color": "green",
                    "size": "small",
                    "x_rel": "0.578334",
                    "y_rel": "0.521314"
                }
            ]
        }
    ]
}
```

]